## **Fertilization and Management Advice**

ANDREA BOLLINI Marsabit Kenya 07991 27346



P.O.Box 1332-00502 Cara House, Karen rd Nairobi/Kenya +254 (0)728 970 136 info@soilcares.com

#### **General Information**

Sample Number: AAASA00737A18 Date: 2018-06-08 Field Name: NURSERY FARM Field Size: 0.03 acre Soil Texture: Sandy Loam

Crop Name : tomato Target Yield : 200 kg

# **Actual Nutrient Need(in kg)**

|           | 0.2 kg   | 0.0 kg | 0.0 kg | 0.0 kg | 23.4 kg        |
|-----------|----------|--------|--------|--------|----------------|
| Parameter | Nitrogen | Sulfur | Iron   | Mn     | organic matter |

### **Fertilizer Recommendations**

Activities Instructions Best Option First Alternative Second Alternative

----- If Available 24 kg Compost or Animal

Manu

Before Planting

Place the fertiliser at the bottom of the planting holes, put 10 cm of

soil on top, add the seed and cover the seed with soil.

1 kg Mav 30:5:5 & 5M and 1 kg CAN 1 kg Mav 30:5:5 & 5M and 1 kg urea 1 kg Mav 30:5:5 & 5M and 1 kg Mav 26:0:0

7

2

3

Y

You can topdress when your crops are 6 weeks old and healthy (no pests, sufficient rain).

6 weeks after

## **Suitable Crop Types**

Potatoes Beans Grains Vegetables









Your soil is suitable for growing potatoes, grains, vegetables and beans.

### **Fertilization and Management Advice**

ANDREA BOLLINI Marsabit Kenya 07991 27346



P.O.Box 1332-00502 Cara House, Karen rd Nairobi/Kenya +254 (0)728 970 136 info@soilcares.com

#### **General Information**

Sample Number: AAASA00737A18 Date: 2018-06-08 Field Name: NURSERY FARM Field Size: 0.03 acre Soil Texture: Sandy Loam

Crop Name : onion Target Yield : 200 kg

# **Actual Nutrient Need(in kg)**

|           | 0.2 kg   | 0.0 kg | 0.0 kg | 0.0 kg | 23.4 kg        |
|-----------|----------|--------|--------|--------|----------------|
| Parameter | Nitrogen | Sulfur | Zinc   | Mn     | organic matter |

#### **Fertilizer Recommendations**

Activities Instructions Best Option First Alternative Second Alternative

\_\_\_\_\_ If Available 24 kg Compost or Animal

Manure

Before Planting

At Planting

Place the fertiliser at the bottom of the planting holes, put 10 cm of

the planting holes, put 10 cm of soil on top, add the seed and cover the seed with soil.

1 kg Mav 30:5:5 & 5M

1 kg Mav 25:5:5 & 4M and 1 kg urea 1 kg Mav 15:10:18 & 6M and 1 kg urea

3

2

6 weeks after

You can topdress when your crops are 6 weeks old and healthy (no pests, sufficient rain).

## **Suitable Crop Types**

Potatoes Beans Grains Vegetables









Your soil is suitable for growing potatoes, grains, vegetables and beans.

### **Fertilization and Management Advice**

ANDREA BOLLINI Marsabit Kenva 07991 27346



P.O.Box 1332-00502 Cara House, Karen rd Nairobi/Kenya +254 (0)728 970 136 info@soilcares.com

#### **General Information**

Date: 2018-06-08 Field Name: NURSERY FARM Field Size: 0.03 acre Sample Number: AAASA00737A18 Soil Texture: Sandy Loam

Crop Name: carrot Target Yield: 200 kg

# **Actual Nutrient Need(in kg)**

|           | 0.2 kg   | 0.0 kg | 0.0 kg | 23.4 kg        |
|-----------|----------|--------|--------|----------------|
| Parameter | Nitrogen | Sulfur | Mn     | organic matter |

#### **Fertilizer Recommendations**

Activities Instructions Best Option First Aternative Second Atemative

24 kg Compost or Animal If Available

**Before Planting** 

At Planting

2

Place the fertiliser at the bottom of the planting holes, put 10 cm of soil on top, add the seed and cover the seed with soil.

1 kg Mav 30:5:5 & 5M

1 kg May 30:5:5 & 5M and 1 kg CAN

1 kg Mav 30:5:5 & 5M and 1 kg May 26:0:0

# **Suitable Crop Types**

Potatoes Beans Grains Vegetables









Your soil is suitable for growing potatoes, grains, vegetables and beans.

### **Soil Status**

| Parameter                | Unit     | Analysis Result | Range Low | Range High | Low | Adequate | High |
|--------------------------|----------|-----------------|-----------|------------|-----|----------|------|
| pH (KCI)                 | pH Value | 7.2             | 4.90      | 6.40       |     |          | •    |
| Organic Carbon           | g/kg     | 11.7            | 20.00     | 50.00      | •   |          |      |
| Total Nitrogen           | g/kg     | 1.3             | 1.00      | 2.00       |     |          |      |
| Total Phosphorus         | g/kg     | 1.2             | 0.20      | 0.60       |     |          | •    |
| Total Sulfur             | g/kg     | 0.2             | 0.30      | 0.50       | •   |          |      |
| Potassium (exch.)        | mmol+/kg | 8.3             | 1.50      | 3.00       |     |          | •    |
| Calcium (exch.)          | mmol+/kg | 70.0            | 15.00     | 25.00      |     |          | •    |
| Magnesium (exch.)        | mmol+/kg | 13.3            | 4.50      | 10.00      |     |          | •    |
| Zinc (M3)                | mg/kg    | 3.5             | 2.50      | 4.00       |     | •        |      |
| Copper (M3)              | mg/kg    | 0.8             | 1.00      | 2.00       | •   |          |      |
| Cation Exchange Capacity | mmol+/kg | 99.5            | 75.00     | 200.00     |     |          |      |
| Clay                     | %        | 7.3             | 25.00     | 50.00      | •   |          |      |
| Sand                     | %        | 73.1            | 35.00     | 55.00      |     |          | •    |
| Total Aluminium          | g/kg     | 66.4            | 56.00     | 91.00      |     |          |      |
| Total Potassium          | g/kg     | 30.9            | 9.80      | 22.00      |     |          | •    |
| Total Silicium           | g/kg     | 334.8           | 250.00    | 330.00     |     |          | •    |
| Total Iron               | g/kg     | 11.7            | 27.00     | 72.00      |     |          |      |
| Phosphorus (M3)          | mg P/kg  | 285.0           | 20.00     | 40.00      |     |          | •    |
| Total Manganese          | g/kg     | 348             | 610.00    | 2300.00    | •   |          |      |

#### Disclaime

The Analysis Report exclusively relates to the sample presented and examined by the scarner. Soil Cares can not werrant that the Analysis Report relates to the source of the sample was not correctly collected. Please note that the recommendations given in the Analysis Report provide indicative rates, that are only valid for the sample presented and based on parameters included in the analysis request, such as crop type, field size, target yield. The sporadic character of samples and the date of the Analysis Report are fundamental in the interpretation of the Analysis Report are fundamental in the interpretation of the Analysis Report are fundamental in the interpretation of the Analysis Report provides on the fundamental in the interpretation of the Analysis Report are fundamental in the interpretation of the Analysis Report are fundamental in the interpretation of the Analysis Report are fundamental in the interpretation of the Analysis Report are fundamental in the interpretation of the Analysis Report analysis Report are fundamental in the interpretation of the interpretation of the sample present and the date of the Analysis Report are fundamental in the interpretation of the Analysis Report analysis Report are fundamental in the interpretation of the sample present and under no circumstances whatescent shall be liable for any special, including but no interpretation of the preparation of the properation of this report, personal data been sert to Soil Cares in the Netherlands. He further consents to his personal data being used for research and marketing purposes. The recipient may at all times required terms of the preparation of the preparation of the personal data being used for research and marketing purposes. The recipient may at all times required terms are present data or demand that his personal data is removed by contacting Soil Cares by email: info@soil cares by email:

Head of Soil Cares Research - Peter van Erp